

# Bacterial Growth in Leukoreduced RBCs During and Following a 3 hr Temperature Excursion



American Red Cross

Wagner SJ, Hapip CA, Abel LA, American Red Cross Holland Lab, Rockville, MD 20855

## Background

It is not uncommon that power interruptions occur at blood centers and hospitals during extreme weather events. When backup generators fail, refrigerated blood warms beyond 6°C. Previous studies have been conducted which evaluate bacterial growth during 30 minutes (the thirty minute rule) or just prior to transfusion (the 4 hour rule).<sup>1,2</sup> However, no study has been conducted directly comparing bacterial growth in leukoreduced RBCs stored with continuous cold storage compared to RBCs stored with a temperature excursion. This study evaluates the levels of several bacteria, some of which are psychrophiles, in refrigerated leukodepleted RBCs during and following a 3-hour temperature excursion towards room temperature.

### References:

1. Transfusion 2013; 53:851-9
2. Vox Sang 2013;105:100-7

## Experimental Design and Methods

Two ABO, Rh identical leukoreduced RBCs were prepared from whole blood collected from consenting donors (Fenwal, Round Lake). The units were pooled, mixed and inoculated on Day 7 with 10-100 CFU/mL (for quantitative plate assay) of *P. fluorescens*, *Y. enterocolita*, *S. marcescens*, *S. liquefaciens*, *E. coli*, *S. aureus*, *E. cloacae* and *S. caprae*. The pool was divided equally into two units and stored at 1-6°C. On Day 14 one of the two units (TEST) was placed on the laboratory bench (20-24°C) for 3 hours, surface temperature recorded, and then returned to 1-6°C storage. The other unit was maintained in refrigerated storage (control). Samples were taken for quantitative plate assay on Day 7, Day 14 pre-excursion (pre), Day 14 post-excursion (post), Day 28 and Day 42. Experiments were repeated 3 times per organism.

## Results

At the end of the 3-hour temperature excursion, the surface temperature of units were  $18.7 \pm 0.8^\circ\text{C}$ . Bacterial results are expressed as  $(\text{CFU/mL}_{\text{TEST}})/(\text{CFU/mL}_{\text{CONTROL}})$  in the table below. No significant differences were observed between the growth of test and control units during storage for *P. fluorescens*, *E. coli*, *S. aureus*, *E. cloacae* or *S. caprae*. On Day 14 post excursion, the bacterial count in test units containing *Y. enterocolitica* declined approximately 4-fold relative to control units; no differences were observed in other timepoints. For *S. marcescens*, significantly greater levels of bacteria (3.2- and 5.4-fold) were observed in test units relative to those of controls on Days 28 and 42, respectively. Bacteria counts of *S. liquefaciens* that were subjected to the temperature excursion were 4.4-fold greater than those post-excursion of control units on Day 14, but this difference did not attain statistical significance.

Organism	$\log(\text{CFU/mL})_{\text{TEST}}/\log(\text{CFU/mL})_{\text{CONTROL}}$				
	Day 7	Day 14 pre	Day 14 post	Day 28	Day 42
<i>P. fluorescens</i>	1.0	0.79 ± 0.40	0.69 ± 0.13	1.10 ± 0.19	0.86 ± 0.14
<i>Y. enterocolitica</i>	1.0	0.79 ± 0.11	0.025 ± 0.033	0.83 ± 0.01	0.76 ± 0.11
<i>S. marcescens</i>	1.0	1.27 ± 0.29	1.03 ± 0.43	3.19 ± 0.15	5.37 ± 0.43
<i>S. liquefaciens</i>	1.0	1.42 ± 0.15	4.38 ± 2.58	1.32 ± 0.29	1.19 ± 0.12
<i>E. coli</i>	1.0	0.94 ± 0.46	1.43 ± 0.43	No growth	No growth
<i>S. aureus</i>	1.0	1.31 ± 0.60	1.00 ± 0.17	0.51 ± 0.11	1.76 ± 0.89
<i>E. cloacae</i>	1.0	0.99 ± 0.25	1.29 ± 0.175	No growth	No growth
<i>S. caprae</i>	1.0	0.33 ± 0.57	0.39 ± 0.35	No growth	No growth

## Conclusions

A 3-hour room temperature excursion in RBCs did not affect bacterial counts in *P. fluorescens*, *E. coli*, *S. aureus*, *E. cloacae*, and *S. caprae*, temporarily decreased the levels of *Y. enterocolitica* immediately following temperature excursion but enhanced the growth of *S. marcescens* during later storage relative to continuous 1-6°C storage.